

Bibliographie de la conférence du 6 avril, dans l'ordre d'apparition des références.

1. Lira A, Angelini R, Le Loc'h F, et al (2018) Trophic flow structure of a neotropical estuary in northeastern Brazil and the comparison of ecosystem model indicators of estuaries. *J Mar Syst* 182:31–45. <https://doi.org/10.1016/j.jmarsys.2018.02.007>
2. Wardeh M, Risley C, McIntyre MK, et al (2015) Database of host-pathogen and related species interactions, and their global distribution. *Sci Data* 2:150049. <https://doi.org/10.1038/sdata.2015.49>
3. Chandra D, Srivastava R, Sharma A (2016) Environment friendly Phosphorus Biofertilizer as an Alternative to Chemical Fertilizers. pp 43–71
4. van der Heijden MG, Bruin S de, Luckerhoff L, et al (2016) A widespread plant-fungal-bacterial symbiosis promotes plant biodiversity, plant nutrition and seedling recruitment. *ISME J* 10:389–399. <https://doi.org/10.1038/ismej.2015.120>
5. Zilber-Rosenberg I, Rosenberg E (2008) Role of microorganisms in the evolution of animals and plants: the hologenome theory of evolution. *FEMS Microbiol Rev* 32:723–735. <https://doi.org/10.1111/j.1574-6976.2008.00123.x>
6. Carrillo J, Ingwell LL, Li X, Kaplan I (2019) Domesticated tomatoes are more vulnerable to negative plant–soil feedbacks than their wild relatives. *J Ecol* 107:1753–1766. <https://doi.org/10.1111/1365-2745.13157>
7. Martín-Robles N, Lehmann A, Seco E, et al (2018) Impacts of domestication on the arbuscular mycorrhizal symbiosis of 27 crop species. *New Phytol* 218:322–334. <https://doi.org/10.1111/nph.14962>
8. York LM, Galindo-Castañeda T, Schussler JR, Lynch JP (2015) Evolution of US maize (*Zea mays* L.) root architectural and anatomical phenes over the past 100 years corresponds to increased tolerance of nitrogen stress. *J Exp Bot* 66:2347–2358. <https://doi.org/10.1093/jxb/erv074>
9. Shi S, Chang J, Tian L, et al (2019) Comparative analysis of the rhizomicrobiome of the wild versus cultivated crop: insights from rice and soybean. *Arch Microbiol* 201:879–888. <https://doi.org/10.1007/s00203-019-01638-8>
10. Abeyasinghe G, Kuchira M, Kudo G, et al (2020) Fungal mycelia and bacterial thiamine establish a mutualistic growth mechanism. *Life Sci Alliance* 3:. <https://doi.org/10.26508/lsa.202000878>
11. Hoffman MT, Arnold AE (2010) Diverse Bacteria Inhabit Living Hyphae of Phylogenetically Diverse Fungal Endophytes. *Appl Environ Microbiol* 76:4063–4075. <https://doi.org/10.1128/AEM.02928-09>
12. Agler MT, Ruhe J, Kroll S, et al (2016) Microbial Hub Taxa Link Host and Abiotic Factors to Plant Microbiome Variation. *PLOS Biol* 14:e1002352. <https://doi.org/10.1371/journal.pbio.1002352>
13. Chelius MK, Triplett EW (2000) Immunolocalization of dinitrogenase reductase produced by *Klebsiella pneumoniae* in association with *Zea mays* L. *Appl Environ Microbiol* 66:783–787. <https://doi.org/10.1128/aem.66.2.783-787.2000>

14. Floc'h J-B, Hamel C, Harker KN, St-Arnaud M (2020) Fungal Communities of the Canola Rhizosphere: Keystone Species and Substantial Between-Year Variation of the Rhizosphere Microbiome. *Microb Ecol.* <https://doi.org/10.1007/s00248-019-01475-8>
15. Floc'h J-B, Hamel C, Lupwayi N, et al (2020) Bacterial Communities of the Canola Rhizosphere: Network Analysis Reveals a Core Bacterium Shaping Microbial Interactions. *Front Microbiol* 11:. <https://doi.org/10.3389/fmicb.2020.01587>
16. Nahar K, Floc'h J-B, Goyer C, et al (2020) Diversity of Soil Bacterial Community Is Influenced by Spatial Location and Time but Not Potato Cultivar. *Phytobiomes J* PBIOMES-01-20-0. <https://doi.org/10.1094/PBIOMES-01-20-0002-R>